




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12Q 1/18, G01N 33/94, C07K 14/315, 14/32		A1	(11) International Publication Number: WO 00/20627
			(43) International Publication Date: 13 April 2000 (13.04.00)
(21) International Application Number: PCT/US99/22665 (22) International Filing Date: 30 September 1999 (30.09.99) (30) Priority Data: 09/163,445 30 September 1998 (30.09.98) US (71) Applicant (for all designated States except US): MILLENNIUM PHARMACEUTICALS, INC. [US/US]; 75 Sidney Street, Cambridge, MA 02139 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): FRITZ, Christian [DE/US]; 14 Brouds Avenue, Natick, MA 01760 (US). YOUNGMAN, Philip [US/US]; 92 Charles Street #53, Boston, MA 02139 (US). GUZMAN, Luz-Maria [MX/US]; 52 Athol Street, Boston, MA 02134 (US). (74) Agents: ELLISON, Eldora, L.; Fish & Richardson P.C., 601 Thirteenth Street, N.W., Washington, DC 20005 (US) et al.		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	

(54) Title: USE OF YNES, ESSENTIAL BACTERIAL GENES AND POLYPEPTIDES

Non-polar gene knockouts in *S. pneumoniae*

MILLENNIUM PHARMACEUTICALS, INC.

(57) Abstract

Disclosed is a gene, termed "S-yneS", found in *Streptococcus pneumoniae*, which is essential for survival for a wide range of bacteria. This gene and the S-yneS polypeptide that it encodes, as well as homologs and orthologs thereof (collectively referred to as "yneS" genes and polypeptides) can be used to identify antibacterial agents for treating a broad spectrum of bacterial infections.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

USE OF YNEs, ESSENTIAL BACTERIAL GENES AND POLYPEPTIDESField of the Invention

The invention relates to the use of yneS, an
10 essential bacterial gene, in identifying antibacterial
agents.

Background of the Invention

Bacterial infections may be cutaneous,
subcutaneous, or systemic. Opportunistic bacterial
15 infections proliferate, especially in patients afflicted
with AIDS or other diseases that compromise the immune
system. Most bacteria that are pathogenic to humans are
gram positive bacteria. The bacterium *Streptococcus*
pneumoniae, for example, typically infects the
20 respiratory tract and can cause lobar pneumonia, as well
as meningitis, sinusitis, and other infections.

Summary of the Invention

The invention is based on the discovery that the
yneS gene of the gram positive bacterium *Streptococcus*
25 *pneumoniae*, termed "S-yneS," and of *Bacillus subtilis*,
termed "B-yneS," are essential for survival. The terms
"yneS gene" and "yneS polypeptide" refer to the S-yneS
and B-yneS genes, as well as their homologs and
orthologs, collectively. While "homologs" are
30 structurally similar genes contained within a species,
"orthologs" are functionally equivalent genes from other
species (within or outside of a given genus). Orthologs

- 2 -

can be identified, for example, in a conventional complementation assay. In addition, such orthologs can be expected to exist in bacteria in the same branch of the phylogenetic tree, as set forth, for example, at
5 ftp://ftp.cme.msu.edu/pub/ RDP/SSU_rRNA/SSU/Prok.phylo.
For example, *B. subtilis* is in the *B. subtilis* subgroup of the *B. subtilis* group in the *Bacillus-Lactobacillus-Streptococcus* Subdivision of the Gram positive phylum. Likewise, *S. pneumoniae* belong to the *Stc. pneumonia*
10 subgroup of *Streptococci*, which also are in the *Bacillus-Lactobacillus-Streptococcus* subdivision of the Gram positive phylum. Other bacteria within the same phylum (particularly, bacteria within the same subdivision, group, or subgroup) can be expected to contain an
15 ortholog of the *yneS* gene described herein, and the ortholog is expected to be essential for survival.

The *yneS* gene is considered an "essential" gene, and the *yneS* polypeptide is considered an "essential" polypeptide, meaning that the function of the gene and
20 polypeptide are necessary for survival of the organism (i.e., *B. subtilis*, *S. pneumoniae*, and other gram positive bacteria). A *yneS* gene and polypeptide can be used in methods for identifying similar genes in pathogenic and non-pathogenic microorganisms. A *yneS*
25 polypeptide can be used to identify compounds that are inhibitors of the pathogens in which the *yneS* polypeptide is expressed and essential. Such inhibitors attenuate bacterial growth by inhibiting the activity of a *yneS* polypeptide, or by inhibiting transcription of a *yneS*
30 gene or translation of the mRNA transcribed from the *yneS* gene.

The coding strand, amino acid sequence, and non-coding strand of *S-yneS* are set forth in Fig. 1 as SEQ ID NOs:1, 2, and 11, respectively. The coding strand, amino

- 3 -

acid, and non-coding strand of B-yneS are set forth in Fig. 2 as SEQ ID NOs:3, 4, and 12 respectively.

Since, as described herein, the yneS gene has been identified and shown to be essential for survival of gram positive bacteria such as *S. pneumoniae* and *B. subtilis*, the yneS gene and polypeptide (including homologs and orthologs of the sequences disclosed herein) can be used to identify antibacterial agents. Such antibacterial agents can readily be identified with high throughput assays to detect inhibition of the metabolic pathway in which the yneS polypeptide participates. This inhibition can be caused by small molecules interacting with (e.g., binding directly or indirectly to) the yneS polypeptide or other essential polypeptides in that pathway.

In an exemplary assay, but not the only assay, a promoter that responds to depletion of the essential yneS polypeptide by upregulation or downregulation is linked to a reporter gene. To identify a promoter that is up- or down-regulated by the depletion of a yneS protein, the gene encoding the yneS protein is deleted from the genome and replaced with a version of the gene in which the sequence encoding the yneS protein is operably linked to a regulatable promoter. The cells containing this regulatable genetic construct are kept alive by the essential yneS polypeptide produced from the genetic construct containing the regulatable promoter. However, the regulatable promoter allows the expression of the yneS polypeptide to be reduced to a level that causes growth inhibition. Total RNA prepared from bacteria under such growth-limiting conditions is compared with RNA from wild-type cells. Standard methods of transcriptional profiling can be used to identify mRNA species that are either more or less abundant (i.e., up- or down-regulated) when expressed under the limiting conditions. Genomic sequence information, e.g., from

- 4 -

GenBank, can be used to identify a promoter that drives expression of the identified RNA species. Such promoters are up- or down-regulated by depletion of the yneS polypeptide.

5 Having identified a promoter(s) that is up- or down-regulated by depletion of the yneS polypeptide, the promoter(s) is operably linked to a reporter gene (e.g., β -galactosidase, gus, or green fluorescent protein (GFP)). A bacterial strain containing this reporter gene
10 construct is then exposed to test compounds. Compounds that inhibit the yneS polypeptide (or other polypeptides in the essential pathway in which the yneS polypeptide participates) will cause a functional depletion of the yneS polypeptide and therefore lead to an upregulation or
15 downregulation of expression of the reporter gene. Because the polypeptides described herein are essential for the survival of bacteria, such as gram positive bacteria, compounds that inhibit the yneS polypeptides in such an assay are expected to be antibacterial and can be
20 further tested, if desired, in standard susceptibility assays.

Another suitable method for identifying antibacterial compounds involves screening for small molecules that specifically interact with (i.e., bind
25 directly or indirectly to) the essential yneS polypeptide. A variety of suitable interaction and binding assays are known in the art as described, for example, in U.S. Patent Nos. 5,585,277 and 5,679,582, incorporated herein by reference. For example, in
30 various conventional assays, test compounds can be assayed for their ability to interact with a yneS polypeptide by measuring the ability of the small molecule to stabilize the yneS polypeptide in its folded, rather than unfolded, state. More specifically, one can
35 measure the degree of protection against unfolding that

- 5 -

is afforded by the test compound. Test compounds that bind the yneS polypeptide with high affinity cause, for example, a large shift in the temperature at which the polypeptide is denatured. Test compounds that stabilize
5 the yneS polypeptide in a folded state can be further tested for antibacterial activity in a standard susceptibility assay.

In a related method for identifying antibacterial compounds, the essential yneS polypeptide is used to
10 isolate peptide or nucleic acid ligands that specifically bind the yneS polypeptide. These peptide or nucleic acid ligands are then used in a displacement screen to identify small molecules that interact with the yneS polypeptide. Such assays can be carried out essentially
15 as described above.

Another suitable method for identifying inhibitors of the essential yneS polypeptide involves identifying a biochemical activity of the polypeptide and then screening for small molecule inhibitors of the activity
20 using, for example, a high throughput screening method.

The various yneS polypeptides can be used, separately or together, in assays to identify test compounds that interact with these polypeptides. Test compounds that interact with these polypeptides then can
25 readily be tested, in conventional assays, for their ability to inhibit bacterial growth. Test compounds that interact with the yneS polypeptides are candidate antibacterial agents, in contrast to compounds that do not interact with the yneS polypeptides. As described
30 herein, any of a variety of art-known methods can be used to assay for the interaction of test compounds with the yneS polypeptides.

The invention also includes a method for identifying an antibacterial agent where the method
35 entails: (a) contacting an essential yneS polypeptide

- 6 -

with a test compound; (b) detecting binding of the test compound to the polypeptide; and, optionally, (c) determining whether a test compound that binds to the yneS polypeptide inhibits growth of bacteria, relative to growth of bacteria cultured in the absence of the test compound that binds to the yneS polypeptide, as an indication that the test compound is an antibacterial agent.

In still another method, interaction of a test compound with a yneS polypeptide (e.g., binding) can be detected in a conventional two-hybrid system for detecting protein/protein interactions (e.g., in yeast or mammalian cells). A test compound found to interact with the yneS polypeptide can be further tested for antibacterial activity in a conventional susceptibility assay. Generally, in such two-hybrid methods, (a) the yneS polypeptide is provided as a fusion protein that includes the polypeptide fused to (i) a transcription activation domain of a transcription factor or (ii) a DNA-binding domain of a transcription factor; (b) the test polypeptide is provided as a fusion protein that includes the test polypeptide fused to (i) a transcription activation domain of a transcription factor or (ii) a DNA-binding domain of a transcription factor; and (c) binding of the test polypeptide to the polypeptide is detected as a reconstitution of a transcription factor. Homologs and orthologs of S-yneS and B-yneS polypeptide described herein can be used in such methods. Reconstitution of the transcription factor can be detected, for example, by detecting transcription of a gene that is operably linked to a DNA sequence bound by the DNA-binding domain of the reconstituted transcription factor (See, for example, White, 1996, Proc. Natl. Acad. Sci. 93:10001-10003 and references

- 7 -

cited therein and Vidal et al., 1996, Proc. Natl. Acad. Sci. 93:10315-10320).

In an alternative method, an isolated nucleic acid molecule encoding a yneS polypeptide is used to identify
5 a compound that decreases the expression of a yneS polypeptide *in vivo* (i.e., in a cell). Such compounds can be used as antibacterial agents. To discover such compounds, cells that express a yneS polypeptide are cultured, exposed to a test compound (or a mixture of
10 test compounds), and the level of expression or activity is compared with the level of yneS polypeptide expression or activity in cells that are otherwise identical but that have not been exposed to the test compound(s). Many standard quantitative assays of gene expression can be
15 utilized in this aspect of the invention.

To identify compounds that modulate expression of a yneS polypeptide, the test compound(s) can be added at varying concentrations to the culture medium of cells that express a yneS polypeptide, as described herein.
20 Such test compounds can include small molecules (typically, non-protein, non-polysaccharide chemical entities), polypeptides, and nucleic acids. The expression of the yneS polypeptide is then measured, for example, by Northern blot PCR analysis or RNase
25 protection analyses using a nucleic acid molecule described herein as a probe. The level of expression in the presence of the test molecule, compared with the level of expression in its absence, will indicate whether or not the test molecule alters the expression of the
30 yneS polypeptide. Because the yneS polypeptides (including homologs and orthologs of the sequences disclosed herein) are essential for survival, test compounds that inhibit the expression and/or function of the yneS polypeptide will inhibit the growth of, or kill,
35 cells that express such yneS polypeptides.

- 8 -

Test compounds can also be assayed for their ability to inhibit cell growth in an overexpression/underexpression screen. In such an assay, the wild-type yneS gene in a cell is replaced with a yneS gene that is under the control of a regulated promoter. By adjusting the level of inducer added to the cell culture, one can produce (i) a cell that expresses more yneS protein than does the wild-type cell (i.e., overexpression) and (ii) a cell that expresses less yneS protein than does the wild-type cell (i.e., underexpression). By adjusting the level of inducer, the level of yneS expression can be increased or decreased (e.g., by at least 25%, 50%, 100%, 200%, or even 500% or greater). Optimally, the inducer is added at low levels to the cells used for underexpression, such that the cells are barely able to grow. Such cells are growth-limited by yneS protein activity, and should therefore be especially sensitive to yneS inhibitors. The cells used for underexpression of yneS can be used to detect inhibitors. A comparison of the effects of test compounds on the cells used for underexpression with the effects on the cells used for overexpression, or with the wild-type strain, which should be more resistant than the cells used for underexpression, provides an indication of the specificity of the test compound.

Typically, the test compound will be a small organic molecule. Alternatively, the test compound can be a test polypeptide (e.g., a polypeptide having a random or predetermined amino acid sequence; or a naturally-occurring or synthetic polypeptide) or a nucleic acid, such as a DNA or RNA molecule. The test compound can be a naturally-occurring compound or it can be synthetically produced, if desired. Synthetic libraries, chemical libraries, and the like can be screened to identify compounds that bind the yneS

- 9 -

polypeptide. More generally, binding of test a compound to the yneS polypeptide can be detected either *in vitro* or *in vivo*. If desired, the above-described methods for identifying compounds that modulate the expression of yneS polypeptides can be combined with measuring the levels of the yneS polypeptides expressed in the cells, e.g., by performing a Western blot analysis using antibodies that bind a yneS polypeptide.

Regardless of the source of the test compound, the yneS polypeptides described herein can be used to identify compounds that inhibit the activity of a yneS protein or transcription of a yneS gene or translation of the mRNA transcribed from the yneS gene. These antibacterial agents can be used to inhibit a wide spectrum of pathogenic or non-pathogenic bacterial strains, particularly gram positive bacteria.

In other embodiments, the invention includes pharmaceutical formulations that include a pharmaceutically acceptable excipient and an antibacterial agent identified using the methods described herein. In particular, the invention includes pharmaceutical formulations that contain antibacterial agents that inhibit the growth of, or kill, pathogenic bacterial strains (e.g., pathogenic gram positive bacterial strains such as pathogenic *Streptococcus* strains). Such pharmaceutical formulations can be used in a method of treating a bacterial infection in an organism (e.g., a *Streptococcus* infection). Such a method entails administering to the organism a therapeutically effective amount of the pharmaceutical formulation, i.e., an amount sufficient to ameliorate signs and/or symptoms of the bacterial infection. In particular, such pharmaceutical formulations can be used to treat bacterial infections in mammals such as humans and domesticated mammals (e.g., cows, pigs, dogs, and

- 10 -

cats), and in plants. The efficacy of such antibacterial agents in humans can be estimated in an animal model system well known to those of skill in the art (e.g., mouse and rabbit model systems of, for example, streptococcal pneumonia).

Various affinity reagents that are permeable to the microbial membrane (i.e., antibodies and antibody fragments) are useful in practicing the methods of the invention. For example polyclonal and monoclonal antibodies that specifically bind to the yneS polypeptide can facilitate detection of yneS polypeptides in various bacterial strains (or extracts thereof). These antibodies also are useful for detecting binding of a test compound to yneS polypeptides (e.g., using the assays described herein). In addition, monoclonal antibodies that bind yneS polypeptides can themselves be used as antibacterial agents.

The invention further features methods of identifying from a large group of mutant strains those strains that have conditional lethal mutations. In general, the gene and corresponding gene product are subsequently identified, although the strains themselves can be used in screening or diagnostic assays. The mechanism(s) of action for the identified genes and gene products provide a rational basis for the design of antibacterial therapeutic agents. These antibacterial agents reduce the action of the gene product in a wild type strain, and therefore are useful in treating a subject with that type, or a similarly susceptible type, of infection by administering the agent to the subject in a pharmaceutically effective amount. Reduction in the action of the gene product includes competitive inhibition of the gene product for the active site of an enzyme or receptor; non-competitive inhibition; disrupting an intracellular cascade path which requires

- 11 -

the gene product; binding to the gene product itself, before or after post-translational processing; and acting as a gene product mimetic, thereby down-regulating the activity. Therapeutic agents include monoclonal
5 antibodies raised against the gene product.

Furthermore, the presence of the gene sequence in certain cells (e.g., a pathogenic bacterium of the same genus or similar species), and the absence or divergence of the sequence in host cells can be determined, if
10 desired. Therapeutic agents directed toward genes or gene products that are not present in the host have several advantages, including fewer side effects, and lower overall dosage.

In various embodiments, the yneS polypeptide used
15 in the assays described herein is derived from a non-pathogenic or pathogenic gram positive bacterium. For example, a yneS polypeptide can be derived from a *Streptococcus* strain, such as *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Streptococcus agalactiae*,
20 *Streptococcus endocarditis*, *Streptococcus faecium*, *Streptococcus sanguis*, *Streptococcus viridans*, and *Streptococcus hemolyticus*. Suitable orthologs of the S-yneS gene can be derived from a wide spectrum of bacteria, such as *Bacillus subtilis*.

25 The invention offers several advantages. For example, the methods for identifying antibacterial agents can be configured for high throughput screening of numerous candidate antibacterial agents. Because the essential yneS gene disclosed herein is thought to be
30 highly conserved, antibacterial drugs targeted to this gene or its gene products are expected to have a broad spectrum of antibacterial activity.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as
35 commonly understood by one of ordinary skill in the art

- 12 -

to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described
5 herein. All publications, patent applications, patents, and other references mentioned herein are incorporated herein by reference in their entirety. In the case of a conflict, the present specification, including definitions, will control. In addition, the materials,
10 methods, and examples are illustrative and are not intended to limit the scope of the invention, which is defined by the claims.

Other features and advantages of the invention will be apparent from the following detailed description,
15 and from the claims.

Brief Description of the Drawings

Fig. 1 is a listing of the nucleic acid and amino acid sequences of a *yneS* gene and polypeptide from a *Streptococcus pneumoniae* strain (SEQ ID NOs:1 and 2,
20 respectively). The complement of the nucleic acid sequence is set forth as SEQ ID NO:11.

Fig. 2 is a listing of the nucleic acid and amino acid sequences of a *yneS* gene and polypeptide from a *B. subtilis* strain (SEQ ID NOs:3 and 4, respectively).
25 The complement of the nucleic acid sequence is set forth as SEQ ID NO:12.

Fig. 3 is a schematic representation of the PCR strategy used to produce DNA molecules used for targeted deletions of essential genes in *Streptococcus pneumoniae*.

30 Fig. 4 is a schematic representation of the strategy used to produce targeted deletions of essential genes in *Streptococcus pneumoniae*.

- 13 -

Detailed Description of the Invention

A gene in the bacterium *Streptococcus pneumoniae* has been found to be essential for the survival of this bacterium. This so-called essential gene, S-yneS, encodes what is referred to herein as a S-yneS polypeptide. The S-yneS gene can be used to identify similar genes in pathogenic microorganisms, such as pathogenic gram positive bacteria (e.g., *Bacillus*), and a yneS gene from *Bacillus subtilis*, termed B-yneS, has been found to be essential for survival. These yneS polypeptides can be used in methods for identifying compounds that are inhibitors of the pathogens in which yneS polypeptides are expressed (e.g., pathogenic and non-pathogenic bacteria, and gram positive bacteria in particular).

Nucleic acids used in practicing the methods of the invention include both RNA and DNA, including genomic DNA and synthetic (e.g., chemically synthesized) DNA. Nucleic acids can be double-stranded or single-stranded. Where single-stranded, the nucleic acid may be a sense strand or an antisense strand. Nucleic acids can be synthesized using oligonucleotide analogs or derivatives (e.g., inosine or phosphorothioate nucleotides). Such oligonucleotides can be used, for example, to prepare nucleic acids that have altered base-pairing abilities or increased resistance to nucleases.

An isolated nucleic acid is a DNA or RNA that is not immediately contiguous with both of the coding sequences with which it is immediately contiguous (one on the 5' end and one on the 3' end) in the naturally occurring genome of the organism from which it is derived. Thus, in one embodiment, an isolated nucleic acid includes some or all of the 5' non-coding (e.g., promoter) sequences that are immediately contiguous to the coding sequence. The term therefore includes, for

- 14 -

example, a recombinant DNA that is incorporated into a vector, into an autonomously replicating plasmid or virus, or into the genomic DNA of a prokaryote or eukaryote, or which exists as a separate molecule (e.g.,
5 a genomic DNA fragment produced by PCR or restriction endonuclease treatment) independent of other sequences. It also includes a recombinant DNA that is part of a hybrid gene encoding an additional polypeptide sequence. The term "isolated" can refer to a nucleic acid or
10 polypeptide that is substantially free of cellular material, viral material, or culture medium (when produced by recombinant DNA techniques), or chemical precursors or other chemicals (when chemically synthesized). Moreover, an isolated nucleic acid
15 fragment is a nucleic acid fragment that is not naturally occurring as a fragment and would not be found in the natural state.

A nucleic acid sequence that is substantially identical to an S-yneS or B-yneS nucleotide sequence is
20 at least 80% identical to the nucleotide sequence represented by SEQ ID NO:1, as depicted in Fig. 1 or SEQ ID NO:3, as depicted in Fig. 2, respectively. For purposes of comparison of nucleic acids, the length of the reference nucleic acid sequence will generally be at
25 least 40 nucleotides, e.g., at least 60 nucleotides or more nucleotides.

To determine the percent identity of two amino acid sequences or of two nucleic acids, the sequences are aligned for optimal comparison purposes (e.g., gaps can
30 be introduced in the sequence of a first amino acid or nucleic acid sequence for optimal alignment with a second amino or nucleic acid sequence). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position
35 in the first sequence is occupied by the same amino acid

- 15 -

residue or nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical
5 positions shared by the sequences (i.e., % identity = # of identical positions/total # of overlapping positions x 100). Preferably, the two sequences are the same length.

The determination of percent identity or homology between two sequences can be accomplished using a
10 mathematical algorithm. A suitable, mathematical algorithm utilized for the comparison of two sequences is the algorithm of Karlin and Altschul (1990) *Proc. Nat'l Acad. Sci. USA* 87:2264-2268, modified as in Karlin and Altschul (1993) *Proc. Nat'l Acad. Sci. USA* 90:5873-5877.
15 Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul, et al. (1990) *J. Mol. Biol.* 215:403-410. BLAST nucleotide searches can be performed with the NBLAST program, score = 100, wordlength = 12 to obtain nucleotide sequences homologous to yneS nucleic
20 acid molecules described herein. BLAST protein searches can be performed with the XBLAST program, score = 50, wordlength = 3 to obtain amino acid sequences homologous to yneS protein molecules. To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as
25 described in Altschul et al., (1997) *Nucleic Acids Res.* 25:3389-3402. When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (e.g., XBLAST and NBLAST) can be used. See <http://www.ncbi.nlm.nih.gov>. Another example of a
30 mathematical algorithm is the algorithm of Myers and Miller, CABIOS (1989). Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a

- 16 -

PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

The percent identity between two sequences can be determined using techniques similar to those described
5 above, with or without allowing gaps. In calculating percent identity, only exact matches are counted.

The yneS polypeptides useful in practicing the invention include, but are not limited to, recombinant polypeptides and natural polypeptides. Also useful in
10 the invention are nucleic acid sequences that encode forms of yneS polypeptides in which naturally occurring amino acid sequences are altered or deleted. Preferred nucleic acids encode polypeptides that are soluble under normal physiological conditions. Also useful are nucleic
15 acids encoding fusion proteins in which a portion of a yneS polypeptide is fused to an unrelated polypeptide (e.g., a marker polypeptide or a fusion partner) to create a fusion protein. For example, the polypeptide can be fused to a hexa-histidine tag to facilitate
20 purification of bacterially expressed polypeptides, or to a hemagglutinin tag to facilitate purification of polypeptides expressed in eukaryotic cells. Also useful in practicing the methods of the invention are isolated polypeptides (and the nucleic acids that encode these
25 polypeptides) that include a first portion and a second portion; the first portion includes, e.g., a yneS polypeptide, and the second portion includes an immunoglobulin constant (Fc) region or a detectable marker.

30 The fusion partner can be, for example, a polypeptide which facilitates secretion, e.g., a secretory sequence. Such a fused polypeptide is typically referred to as a preprotein. The secretory sequence can be cleaved by the host cell to form the
35 mature protein. Additional useful nucleic acids include

- 17 -

nucleic acids that encode a yneS polypeptide fused to a polypeptide sequence to produce an inactive preprotein. Preproteins can be converted into the active form of the protein by removal of the inactivating sequence.

5 Nucleic acids that hybridize, e.g., under moderate or highly stringent hybridization conditions (as defined herein) to all or a portion of the nucleotide sequences represented by SEQ ID NOs:1 or 3, or their complements, can be used to identify additional homologs or orthologs
10 of S-yneS and B-yneS. The hybridizing portion of the hybridizing nucleic acids is typically at least 15 (e.g., 20, 25, 30, or 50) nucleotides in length. The hybridizing portion of the hybridizing nucleic acid is at least 80%, e.g., at least 95%, or at least 98%, identical
15 to the sequence of a portion or all of a nucleic acid encoding a yneS polypeptide or its complement. Hybridizing nucleic acids of the type described herein can be used as a cloning probe, a primer (e.g., a PCR primer), or a diagnostic probe. Nucleic acids that
20 hybridize to the nucleotide sequences represented by SEQ ID NOs:1 or 3 are considered "antisense oligonucleotides."

Cells that may be used in practicing the methods of the invention include various engineered cells, e.g.,
25 transformed host cells, that contain a yneS nucleic acid described herein. A transformed cell is a cell into which (or into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a nucleic acid encoding a yneS polypeptide. Both prokaryotic and
30 eukaryotic cells are included, e.g., bacteria, such as *Streptococcus*, *Bacillus*, and the like.

Various genetic constructs (e.g., vectors and plasmids) that include a yneS nucleic acid that is operably linked to a transcription and/or translation
35 sequence to enable expression (e.g., expression vectors)

- 18 -

can be used in practicing the methods of the invention. A selected nucleic acid, e.g., a DNA molecule encoding a yneS polypeptide, is "operably linked" to a transcription and/or translation sequence when it is positioned
5 adjacent to one or more sequence elements, e.g., a promoter, which direct transcription and/or translation of the sequence such that the sequence elements can control transcription and/or translation of the selected nucleic acid.

10 The terms "protein" and "polypeptide" both refer to any chain of amino acids, regardless of length or post-translational modification (e.g., glycosylation or phosphorylation). Thus, the term yneS polypeptide includes full-length, naturally occurring, isolated yneS
15 proteins, as well as recombinantly or synthetically produced polypeptides that correspond to the full-length, naturally occurring protein, or to a portion of the naturally occurring or synthetic polypeptide.

A purified or isolated compound is a composition
20 that is at least 60% by weight the compound of interest, e.g., a yneS polypeptide or antibody. Preferably the preparation is at least 75% (e.g., at least 90%, 95%, or even 99%) by weight the compound of interest. Purity can be measured by any appropriate standard method, e.g.,
25 column chromatography, polyacrylamide gel electrophoresis, or HPLC analysis.

Preferred yneS polypeptides include a sequence substantially identical to all or a portion of a naturally occurring yneS polypeptide, e.g., including all
30 or a portion of the sequence shown in Figs. 1 or 2. Polypeptides "substantially identical" to the yneS polypeptide sequence described herein have an amino acid sequence that is at least 80% identical to the amino acid sequence of the yneS polypeptide represented by SEQ ID
35 NO:2 or SEQ ID NO:4 (measured as described herein). The

- 19 -

new polypeptides can also have a greater percentage identity, e.g., 85%, 90%, 95%, or even higher. For purposes of comparison, the length of the reference yneS polypeptide sequence will generally be at least 16 amino acids, e.g., at least 20 or 25 amino acids.

In the case of polypeptide sequences that are less than 100% identical to a reference sequence, the non-identical positions are preferably, but not necessarily, conservative substitutions for the reference sequence. Conservative substitutions typically include substitutions within the following groups: glycine and alanine; valine, isoleucine, and leucine; aspartic acid and glutamic acid; asparagine and glutamine; serine and threonine; lysine and arginine; and phenylalanine and tyrosine.

Where a particular polypeptide is said to have a specific percent identity to a reference polypeptide of a defined length, the percent identity is relative to the reference polypeptide. Thus, a polypeptide that is 50% identical to a reference polypeptide that is 100 amino acids long can be a 50 amino acid polypeptide that is completely identical to a 50 amino acid long portion of the reference polypeptide. Alternatively, it can be a 100 amino acid long polypeptide that is 50% identical to the reference polypeptide over its entire length. Of course, other polypeptides also will meet the same criteria.

The methods of the invention also make use of purified or isolated antibodies that specifically bind to a yneS polypeptide. An antibody "specifically binds" to a particular antigen, e.g., a yneS polypeptide, when it binds to that antigen, but does not substantially recognize and bind to other molecules in a sample, e.g., a biological sample, that naturally includes a yneS polypeptide.

- 20 -

The various nucleic acids described herein can be used in a method of obtaining a gene related to a *yneS* gene. Such a method entails obtaining a labeled probe that includes an isolated nucleic acid which encodes all or a portion of a *yneS* nucleic acid, or a homolog thereof; screening a nucleic acid fragment library with the labeled probe under conditions that allow hybridization of the probe to nucleic acid fragments in the library, thereby forming nucleic acid duplexes; isolating labeled duplexes, if any; and preparing a full-length gene sequence from the nucleic acid fragments in any labeled duplex to obtain a gene related to the *yneS* gene (e.g., homologs and orthologs of *yneS*).

Identifying the *Streptococcus* S-*yneS* Gene

As shown by the experiments described below, the S-*yneS* gene is essential for survival of *Streptococcus pneumoniae*. *Streptococcus pneumoniae* is available from the ATCC. In general, and for the examples set forth below, *yneS* genes can be identified by creating targeted deletions of genes of interest in a bacterium, e.g., *S. pneumoniae*. The gene of interest was selected as follows. Using standard molecular biology techniques, a library containing fragments of the *Streptococcus pneumoniae* genome was made, using M13 phage or plasmid DNA as the vector. Open reading frames (ORFs) contained within this library were randomly sequenced, using primers that hybridized to the vector. The gene of interest selected for targeted deletion satisfied four criteria, as determined by comparing the sequences with the GenBank database of nucleotide sequences: (i) the ORF had no known function; (ii) the ORF had an ortholog in *Bacillus subtilis*; (iii) the ORF was conserved in other bacteria, with $p < 10^{-10}$; and (iv) the ORF had no eukaryotic ortholog, with $p > 10^{-3}$. The *Streptococcus*

- 21 -

gene S-yneS met each of these criteria, suggesting that a compound that inhibited the S-yneS gene or gene product would have a broad spectrum of antibacterial activity.

The S-yneS gene was replaced with a nucleic acid sequence conferring resistance to the antibiotic erythromycin (an "erm" gene). Other genetic markers can be used in lieu of this particular antibiotic resistance marker. Polymerase chain reaction (PCR) amplification was used to make a targeted deletion in the *Streptococcus* genomic DNA, as shown in Fig. 3. Several PCR reactions were used to produce the DNA molecules needed to carry out target deletion of the genes of interest. First, using primers 5 and 6, an erm gene was amplified from pIL252 from *B. subtilis* (available from the *Bacillus* Genetic Stock Center, Columbus, OH). Primer 5 consists of 21 nucleotides that are identical to the promoter region of the erm gene and complementary to Sequence A. Primer 5 has the sequence 5'GTG TTC GTG CTG ACT TGC ACC3' (SEQ ID NO:5). Primer 6 consists of 21 nucleotides that are complementary to the 3' end of the erm gene. Primer 6 has the sequence 5'GAA TTA TTT CCT CCC GTT AAA3' (SEQ ID NO:6). PCR amplification of the erm gene was carried out under the following conditions: 30 cycles of 94°C for 1 minute, 55°C for 1 minute, and 72°C for 1.5 minutes, followed by one cycle of 72°C for 10 minutes.

In the second and third PCR reactions, sequences flanking the gene of interest were amplified and produced as hybrid DNA molecules that also contained a portion of the erm gene. The second reaction produced a double-stranded DNA molecule (termed "Left Flanking Molecule") that includes sequences upstream of the 5' end of the gene of interest and the first 21 nucleotides of the erm gene. As shown in Fig. 3, this reaction utilized primer 1, which is 21 nucleotides in length and identical to a sequence that is located approximately 500 bp upstream of

- 22 -

the translation start site of the gene of interest. Primers 1 and 2 are gene-specific and include the sequences 5'ACC CAC TCC GTG AAG TCC ACC3' (SEQ ID NO:7) and 5'GAA CCC AGC AGA TAG GCT AGG3' 3' (SEQ ID NO:8), respectively, for *yneS*. Primer 2 is 42 nucleotides in length, with 21 of the nucleotides at the 3' end of the primer being complementary to the 5' end of the sense strand of the gene of interest. The 21 nucleotides at the 5' end of the primer were identical to Sequence A and are therefore complementary to the 5' end of the *erm* gene. Thus, PCR amplification using primers 1 and 2 produced the left flanking DNA molecule, which is a hybrid DNA molecule containing a sequence located upstream of the gene of interest and 21 base pairs of the *erm* gene, as shown in Fig. 3.

The third PCR reaction was similar to the second reaction, but produced the right flanking DNA molecule, shown in Fig. 3. The right flanking DNA molecule contains 21 base pairs of the 3' end of the *erm* gene, a 21 base pair portion of the 3' end of the gene of interest, and sequences downstream of the gene of interest. This right flanking DNA molecule was produced with gene-specific primers 3 and 4. For *S-yneS*, primers 3 and 4 included the sequences 5'GAA CCT AAC CCA TCA AGA TCC3' (SEQ ID NO:9) and 5'CAG TTT TAG ATG GCT TTA ACA GCG3' (SEQ ID NO:10), respectively. Primer 3 is 42 nucleotides; the 21 nucleotides at the 5' end of Primer 3 are identical to Sequence B and therefore are identical to the 3' end of the *erm* gene. The 21 nucleotides at the 3' end of Primer 3 are identical to the 3' end of the gene of interest. Primer 4 is 21 nucleotides in length and is complementary to a sequence located approximately 500 bp downstream of the gene of interest.

PCR amplification of the left and right flanking DNA molecules was carried out, separately, in 50 μ l

- 23 -

reaction mixtures containing: 1 μ l *Streptococcus pneumoniae* (RX1) DNA (0.25 μ g), 2.5 μ l Primer 1 or Primer 4 (10 pmol/ μ l), 2.5 μ l Primer 2 or Primer 3 (20 pmol/ μ l), 1.2 μ l a mixture dNTPs (10 mM each), 37 μ l H₂O, 0.7 μ l Taq polymerase (5 U/ μ l), and 5 μ l 10x Taq polymerase buffer (10 mM Tris, 50 mM KCl, 2.5 mM MgCl₂). The left and right flanking DNA molecules were amplified using the following PCR cycling program: 95°C for 2 minutes; 72°C for 1 minute; 94°C for 30 seconds; 49°C for 30 seconds; 72°C for 1 minute; repeating the 94°C, 49°C, and 72°C incubations 30 times; 72°C for 10 minutes and then stopping the reactions. A 15 μ l aliquot of each reaction mixture then was electrophoresed through a 1.2% low melting point agarose gel in TAE buffer and then stained with ethidium bromide. Fragments containing the amplified left and right flanking DNA molecules were excised from the gel and purified using the QIAQUICK™ gel extraction kit (Qiagen, Inc.) Other art-known methods for amplifying and isolating DNA can be substituted. The flanking left and right DNA fragments were eluted into 30 μ l TE buffer at pH 8.0.

The amplified *erm* gene and left and right flanking DNA molecules were then fused together to produce the fusion product, as shown in Fig. 3. The fusion PCR reaction was carried out in a volume of 50 μ l containing: 2 μ l of each of the left and right flanking DNA molecules and the *erm* gene PCR product; 5 μ l of 10x buffer; 2.5 μ l of Primer 1 (10 pmol/ μ l); 2.5 μ l of Primer 4 (10 pmol/ μ l), 1.2 μ l dNTP mix (10 mM each) 32 μ l H₂O, and 0.7 μ l Taq polymerase. The PCR reaction was carried out using the following cycling program: 95°C for 2 minutes; 72°C for 1 minute; 94°C for 30 seconds, 48°C for 30 seconds; 72°C for 3 minutes; repeat the 94°C, 48°C and 72°C incubations 25 times; 72°C for 10 minutes. After the reaction was stopped, a 12 μ l aliquot of the reaction

- 24 -

mixture was electrophoresed through an agarose gel to confirm the presence of a final product of approximately 2 kb.

A 5 μ l aliquot of the fusion product was used to transform *S. pneumoniae* grown on a medium containing erythromycin in accordance with standard techniques. As shown in Fig. 4, the fusion product and the *S. pneumoniae* genome undergo a homologous recombination event so that the *erm* gene replaces the chromosomal copy of the gene of interest, thereby creating a gene knockout. Disruption of a *yneS* gene results in no growth on a medium containing erythromycin. Using this gene knockout method, the *yneS* gene was identified as being essential for survival.

15 Identification of Homologs and Orthologs of the S-yneS Gene

Having shown that the S-yneS gene is essential for survival of *Streptococcus*, it can be expected that homologs and orthologs of this gene, when present in other gram positive bacteria, are essential for survival of those bacteria as well. The coding sequences of S-yneS was used to search the GenBank database of nucleotide sequences, and an ortholog of the S-yneS sequence was identified in *B. subtilis*. Sequence comparisons were performed using the Basic Local Alignment Search Tool (BLAST) (Altschul et al., J. Mol. Biol. 215:403-410, 1990). The percent sequence identity shared by the S-yneS polypeptides and the *B. subtilis* ortholog, B-yneS, was determined using the GAP program from the Genetics Computer Group (GCG) Wisconsin Sequence Analysis Package (Wisconsin Package Version 9.1; Madison, WI). The default parameters for gap weight (12) and length weight (4) were used.

- 25 -

Typically, yneS polypeptides and their homologs or orthologs share at least 25% (e.g., at least 30%, 35%, or 40%) sequence identity. Typically, the DNA sequences encoding yneS polypeptides and their homologs or orthologs share at least 20% (e.g., at least 30%, 35%, 40% or 45%) sequence identity. Bioinformatics analysis of the S-yneS gene showed that this gene is widely conserved among bacteria. To confirm that the identified ortholog of S-yneS, B-yneS, is essential for survival of *B. subtilis*, the B-yneS gene was deleted from the *B. subtilis* genome. Such a deletion strain has been constructed and does not survive, confirming the essential nature of the polypeptide. The fact that the B-yneS gene also is essential for survival suggests that yneS genes are essential in all gram positive bacteria in which they are present. Therefore, an antibacterial drug targeted to a yneS gene or gene product is expected to have a broad spectrum of antibacterial activity.

Identification of YneS Genes and Polypeptides in

20 Additional

Bacterial Strains and Species

Now that the S-yneS and B-yneS genes have been identified as being essential for survival, these genes, or fragments thereof, can be used to detect homologous or orthologous genes in yet other organisms. In particular, these genes can be used to analyze various pathogenic and non-pathogenic strains of bacteria, particularly gram positive bacteria. Fragments of a nucleic acid (DNA or RNA) encoding a yneS polypeptide (including homologs and orthologs of the sequences listed herein, or sequences complementary thereto) can be used as probes in conventional nucleic acid hybridization assays of nucleic acids from bacteria. For example, nucleic acid probes (which typically are 8-30, or usually 15-20, nucleotides

- 26 -

in length) can be used to detect yneS genes in art-known molecular biology methods, such as Southern blotting, Northern blotting, dot or slot blotting, PCR amplification methods, colony hybridization methods, and the like. Typically, an oligonucleotide probe based on the nucleic acid sequences described herein, or fragment thereof, is labeled and used to screen a genomic library constructed from mRNA obtained from a bacterial strain of interest. A suitable method of labeling involves using polynucleotide kinase to add ^{32}P -labeled ATP to the oligonucleotide used as the probe. This method is well known in the art, as are several other suitable methods (e.g., biotinylation and enzyme labeling).

Hybridization of the oligonucleotide probe to the library, or other nucleic acid sample, typically is performed under moderate to highly stringent conditions. Nucleic acid duplex or hybrid stability is expressed as the melting temperature or T_m , which is the temperature at which a probe dissociates from a target DNA. This melting temperature is used to define the required stringency conditions. If sequences are to be identified that are related and substantially identical to the probe, rather than identical, then it is useful to first establish the lowest temperature at which only homologous hybridization occurs with a particular concentration of salt (e.g., SSC or SSPE). Then, assuming that 1% mismatching results in a 1°C decrease in the T_m , the temperature of the final wash in the hybridization reaction is reduced accordingly (for example, if sequences having $\geq 95\%$ identity with the probe are sought, the final wash temperature is decreased by 5°C). In practice, the change in T_m can be between 0.5° and 1.5°C per 1% mismatch.

High stringency hybridization conditions include, for example, hybridizing at 68°C in 5x SSC/5x Denhardt's

- 27 -

solution/1.0% SDS, or in 0.5 M NaHPO₄ (pH 7.2)/1 mM EDTA/7% SDS, or in 50% formamide/0.25 M NaHPO₄ (pH 7.2)/0.25 M NaCl/1 mM EDTA/7% SDS; and washing in 0.2x SSC/0.1% SDS at room temperature or at 42°C, or in 0.1x
5 SSC/0.1% SDS at 68°C, or in 40 mM NaHPO₄ (pH 7.2)/1 mM EDTA/5% SDS at 50°C, or in 40 mM NaHPO₄ (pH 7.2) 1 mM EDTA/1% SDS at 50°C. Moderately stringent conditions include washing in 3x SSC at 42°C. The parameters of
10 salt concentration and temperature can be varied to achieve the optimal level of identity between the probe and the target nucleic acid. Additional guidance regarding such conditions is readily available in the art, for example, by Sambrook et al., 1989, Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press,
15 N.Y.; and Ausubel et al. (eds.), 1995, Current Protocols in Molecular Biology, (John Wiley & Sons, N.Y.) at Unit 2.10.

In one approach, libraries constructed from pathogenic or non-pathogenic bacterial strains can be
20 screened. For example, such strains can be screened for expression of yneS genes by Northern blot analysis. Upon detection of transcripts of a yneS gene, libraries can be constructed from RNA isolated from the appropriate strain, utilizing standard techniques well known to those
25 of skill in the art. Alternatively, a total genomic DNA library can be screened using a yneS gene probe.

New gene sequences can be isolated, for example, by performing PCR using two degenerate oligonucleotide primer pools designed on the basis of nucleotide
30 sequences within the yneS gene (including homologs and orthologs of the sequences disclosed herein), as described herein. The template for the reaction can be DNA obtained from strains known to express, or suspected of expressing, a yneS allele (including alleles of
35 homologs or orthologs of the sequences disclosed herein).

- 28 -

The PCR product can be subcloned and sequenced to ensure that the amplified sequences represent the sequences of a new yneS nucleic acid sequence.

Synthesis of a yneS polypeptide can readily be
5 accomplished using any of the various art-known
techniques. For example, a yneS polypeptide can be
synthesized chemically *in vitro*, or enzymatically (e.g.,
by *in vitro* transcription and translation).
Alternatively, the gene can be expressed in, and the
10 polypeptide purified from, a cell (e.g., a cultured cell)
by using any of the numerous, available gene expression
systems. For example, the polypeptide antigen can be
produced in a prokaryotic host (e.g., *E. coli* or *B.*
subtilis) or in eukaryotic cells, such as yeast cells or
15 in insect cells (e.g., by using a baculovirus-based
expression vector).

Proteins and polypeptides can also be produced in
plant cells, if desired. For plant cells, viral
expression vectors (e.g., cauliflower mosaic virus and
20 tobacco mosaic virus) and plasmid expression vectors
(e.g., Ti plasmid) are suitable. Such cells are
available from a wide range of sources (e.g., the
American Type Culture Collection, Rockland, MD; also,
see, e.g., Ausubel et al., *Current Protocols in Molecular*
25 *Biology*, John Wiley & Sons, New York, 1994). The optimal
methods of transformation or transfection and the choice
of expression vehicle will depend on the host system
selected. Transformation and transfection methods are
described, e.g., in Ausubel et al., *supra*; expression
30 vehicles may be chosen from those provided, e.g., in
Cloning Vectors: A Laboratory Manual (P.H. Pouwels et
al., 1985, Supp. 1987). The host cells harboring the
expression vehicle can be cultured in conventional
nutrient media, adapted as needed for activation of a

- 29 -

chosen gene, repression of a chosen gene, selection of transformants, or amplification of a chosen gene.

If desired, a yneS polypeptide (including homologs and orthologs of the sequences disclosed herein) can be produced as a fusion protein. For example, the expression vector pUR278 (Ruther et al., *EMBO J.*, 2:1791, 1983) can be used to create lacZ fusion proteins. The art-known pGEX vectors can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can be easily purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. The pGEX vectors are designed to include thrombin or factor Xa protease cleavage sites so that the cloned target gene product can be released from the GST moiety.

In an exemplary expression system, a baculovirus such as *Autographa californica* nuclear polyhedrosis virus (AcNPV), which grows in *Spodoptera frugiperda* cells, can be used as a vector to express foreign genes. A coding sequence encoding a yneS polypeptide can be cloned into a non-essential region (for example the polyhedrin gene) of the viral genome and placed under control of a promoter, e.g., the polyhedrin promoter or an exogenous promoter. Successful insertion of a gene encoding a yneS polypeptide can result in inactivation of the polyhedrin gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat encoded by the polyhedrin gene). These recombinant viruses are then typically used to infect insect cells (e.g., *Spodoptera frugiperda* cells) in which the inserted gene is expressed (see, e.g., Smith et al., *J. Virol.*, 46:584, 1983; Smith, U.S. Patent No. 4,215,051). If desired, mammalian cells can be used in lieu of insect cells, provided that the virus is engineered such that the gene encoding the

- 30 -

desired polypeptide is placed under the control of a promoter that is active in mammalian cells.

In mammalian host cells, a number of viral-based expression systems can be utilized. When an adenovirus
5 is used as an expression vector, the nucleic acid sequence encoding the yneS polypeptide can be ligated to an adenovirus transcription/ translation control complex, e.g., the late promoter and tripartite leader sequence. This chimeric gene can then be inserted into the
10 adenovirus genome by *in vitro* or *in vivo* recombination. Insertion into a non-essential region of the viral genome (e.g., region E1 or E3) will result in a recombinant virus that is viable and capable of expressing a yneS gene product in infected hosts (see, e.g., Logan, Proc.
15 Natl. Acad. Sci. USA, 81:3655, 1984).

Specific initiation signals may be required for efficient translation of inserted nucleic acid sequences. These signals include the ATG initiation codon and adjacent sequences. In general, exogenous translational
20 control signals, including, perhaps, the ATG initiation codon, should be provided. Furthermore, the initiation codon must be in phase with the reading frame of the desired coding sequence to ensure translation of the entire sequence. These exogenous translational control
25 signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements, or transcription terminators (Bittner et al., *Methods in*
30 *Enzymol.*, 153:516, 1987).

YneS polypeptides can be expressed individually or as fusions with a heterologous polypeptide, such as a signal sequence or other polypeptide having a specific cleavage site at the N-and/or C-terminus of the protein
35 or polypeptide. The heterologous signal sequence

- 31 -

selected should be one that is recognized and processed, i.e., cleaved by a signal peptidase, by the host cell in which the fusion protein is expressed.

A host cell can be chosen that modulates the expression of the inserted sequences, or modifies and processes the gene product in a specific, desired fashion. Such modifications and processing (e.g., cleavage) of protein products may facilitate optimal functioning of the protein. Various host cells have characteristic and specific mechanisms for post-translational processing and modification of proteins and gene products. Appropriate cell lines or host systems familiar to those of skill in the art of molecular biology can be chosen to ensure the correct modification and processing of the foreign protein expressed. To this end, eukaryotic host cells that possess the cellular machinery for proper processing of the primary transcript, and phosphorylation of the gene product can be used. Such mammalian host cells include, but are not limited to, CHO, VERO, BHK, HeLa, COS, MDCK, 293, 3T3, WI38, and choroid plexus cell lines.

If desired, the yneS polypeptide (including homologs and orthologs of the sequences described herein) can be produced by a stably-transfected mammalian cell line. A number of vectors suitable for stable transfection of mammalian cells are available to the public, see, e.g., Pouwels et al. (supra); methods for constructing such cell lines are also publicly known, e.g., in Ausubel et al. (supra). In one example, DNA encoding the protein is cloned into an expression vector that includes the dihydrofolate reductase (DHFR) gene. Integration of the plasmid and, therefore, the yneS polypeptide-encoding gene into the host cell chromosome is selected for by including 0.01-300 μ M methotrexate in the cell culture medium (as described in Ausubel et al.,

- 32 -

supra). This dominant selection can be accomplished in most cell types.

Recombinant protein expression can be increased by DHFR-mediated amplification of the transfected gene.

5 Methods for selecting cell lines bearing gene amplifications are described in Ausubel et al. (supra); such methods generally involve extended culture in medium containing gradually increasing levels of methotrexate. DHFR-containing expression vectors commonly used for this
10 purpose include pCVSEII-DHFR and pAdD26SV(A) (described in Ausubel et al., supra).

A number of other selection systems can be used, including but not limited to, herpes simplex virus thymidine kinase genes, hypoxanthine-guanine
15 phosphoribosyl-transferase genes, and adenine phosphoribosyltransferase genes, which can be employed in *tk*, *hgp**rt*, or *ap**rt* cells, respectively. In addition, *gpt*, which confers resistance to mycophenolic acid (Mulligan et al., *Proc. Natl. Acad. Sci. USA*, 78:2072,
20 1981); *neo*, which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al., *J. Mol. Biol.*, 150:1, 1981); and *hygro*, which confers resistance to hygromycin (Santerre et al., *Gene*, 30:147, 1981), can be used.

25 Alternatively, any fusion protein can be readily purified by utilizing an antibody or other molecule that specifically binds the fusion protein being expressed. For example, a system described in Janknecht et al., *Proc. Natl. Acad. Sci. USA*, 88:8972 (1981), allows for
30 the ready purification of non-denatured fusion proteins expressed in human cell lines. In this system, the gene of interest is subcloned into a vaccinia recombination plasmid such that the gene's open reading frame is translationally fused to an amino-terminal tag consisting
35 of six histidine residues. Extracts from cells infected

- 33 -

with recombinant vaccinia virus are loaded onto Ni²⁺ nitriloacetic acid-agarose columns, and histidine-tagged proteins are selectively eluted with imidazole-containing buffers.

5 Alternatively, the yneS polypeptide, or a portion thereof, can be fused to an immunoglobulin Fc domain. Such a fusion protein can be readily purified using a protein A column, for example. Moreover, such fusion proteins permit the production of a chimeric form of a
10 yneS polypeptide having increased stability *in vivo*.

 Once the recombinant yneS polypeptide is expressed, it can be isolated (i.e., purified). Secreted forms of the polypeptides can be isolated from cell culture media, while non-secreted forms must be isolated
15 from the host cells. Polypeptides can be isolated by affinity chromatography. For example, an anti-yneS antibody (e.g., produced as described herein) can be attached to a column and used to isolate the protein. Lysis and fractionation of cells harboring the protein
20 prior to affinity chromatography can be performed by standard methods (see, e.g., Ausubel et al., supra). Alternatively, a fusion protein can be constructed and used to isolate a yneS polypeptide (e.g., a yneS-maltose binding fusion protein, a yneS- β -galactosidase fusion
25 protein, or a yneS-trpE fusion protein; see, e.g., Ausubel et al., supra; New England Biolabs Catalog, Beverly, MA). The recombinant protein can, if desired, be further purified, e.g., by high performance liquid chromatography using standard techniques (see, e.g.,
30 Fisher, *Laboratory Techniques In Biochemistry And Molecular Biology*, eds., Work and Burdon, Elsevier, 1980).

 Given the amino acid sequences described herein, polypeptides useful in practicing the invention,
35 particularly fragments of yneS polypeptides, can be

- 34 -

produced by standard chemical synthesis (e.g., by the methods described in *Solid Phase Peptide Synthesis*, 2nd ed., The Pierce Chemical Co., Rockford, IL, 1984) and used as antigens, for example.

5 Assay for Antibacterial Agents

The invention provides methods for identifying antibacterial agent. Although the inventor is not bound by any particular theory as to the biological mechanism involved, the new antibacterial agents are thought to
10 inhibit specifically (1) the function of a yneS polypeptide or (2) expression of a yneS gene. In preferred methods, screening for antibacterial agents is accomplished by identifying those compounds (e.g., small organic molecules) that inhibit the activity of a yneS
15 polypeptide or the expression of a yneS gene.

In an exemplary assay, but not the only assay, a promoter that responds to depletion of the yneS polypeptide by upregulation or downregulation is linked to a reporter gene (e.g., β -galactosidase, gus, or GFP),
20 as described above. A bacterial strain containing this reporter gene construct is then exposed to test compounds. Compounds that inhibit the yneS polypeptide (or other polypeptides in the essential pathway in which the yneS polypeptide participates) will cause a
25 functional depletion of the yneS polypeptide and therefore lead to an upregulation or downregulation of expression the reporter gene. Because the polypeptides described herein are essential for the survival of bacteria, compounds that inhibit the yneS polypeptides in
30 such an assay are expected to be antibacterial agents and can be further tested, if desired, in conventional susceptibility assays.

In other suitable methods, screening for antibacterial agents is accomplished by (i) identifying

- 35 -

those compounds that interact with or bind to a yneS polypeptide and (ii) further testing such compounds for their ability to inhibit bacterial growth *in vitro* or *in vivo*.

5 Specific binding of a test compound to a polypeptide can be detected, for example, *in vitro* by reversibly or irreversibly immobilizing the test compound(s) on a substrate, e.g., the surface of a well of a 96-well polystyrene microtitre plate. Methods for
10 immobilizing polypeptides and other small molecules are well known in the art. For example, the microtitre plates can be coated with a yneS polypeptide(s) by adding the polypeptide(s) in a solution (typically, at a concentration of 0.05 to 1 mg/ml in a volume of 1-100 μ l)
15 to each well, and incubating the plates at room temperature to 37°C for 0.1 to 36 hours. Polypeptides that are not bound to the plate can be removed by shaking the excess solution from the plate, and then washing the plate (once or repeatedly) with water or a buffer.
20 Typically, the polypeptide is in water or a buffer. The plate is then washed with a buffer that lacks the bound polypeptide. To block the free protein-binding sites on the plates, the plates are blocked with a protein that is unrelated to the bound polypeptide. For example, 300 μ l
25 of bovine serum albumin (BSA) at a concentration of 2 mg/ml in Tris-HCl is suitable. Suitable substrates include those substrates that contain a defined cross-linking chemistry (e.g., plastic substrates, such as polystyrene, styrene, or polypropylene substrates from
30 Corning Costar Corp. (Cambridge, MA), for example). If desired, a beaded particle, e.g., beaded agarose or beaded sepharose, can be used as the substrate.

Interaction of the test compound with yneS polypeptide(s) can be detected by any of a variety of
35 art-known methods. For example, an antibody that

- 36 -

specifically binds a yneS polypeptide can be used in an immunoassay. If desired, the antibody can be labeled (e.g., fluorescently or with a radioisotope) and detected directly (see, e.g., West and McMahon, *J. Cell Biol.* 5 74:264, 1977). Alternatively, a second antibody can be used for detection (e.g., a labeled antibody that binds the Fc portion of an anti-YneS antibody). In an alternative detection method, the yneS polypeptide is labeled, and the label is detected (e.g., by labeling a 10 yneS polypeptide with a radioisotope, fluorophore, chromophore, or the like). In still another method, the yneS polypeptide is produced as a fusion protein with a protein that can be detected optically, e.g., green fluorescent protein (which can be detected under UV 15 light). In an alternative method, the polypeptide (e.g., yneS) can be produced as a fusion protein with an enzyme having a detectable enzymatic activity, such as horse radish peroxidase, alkaline phosphatase, β -galactosidase, or glucose oxidase. Genes encoding all of these enzymes 20 have been cloned and are readily available for use by those of skill in the art. If desired, the fusion protein can include an antigen, and such an antigen can be detected and measured with a polyclonal or monoclonal antibody using conventional methods. Suitable antigens 25 include enzymes (e.g., horse radish peroxidase, alkaline phosphatase, and β -galactosidase) and non-enzymatic polypeptides (e.g., serum proteins, such as BSA and globulins, and milk proteins, such as caseins).

In various cell-based methods for identifying 30 polypeptides that bind yneS polypeptides, the conventional two-hybrid assays of protein/protein interactions can be used (see e.g., Chien et al., *Proc. Natl. Acad. Sci. USA*, 88:9578, 1991; Fields et al., U.S. Pat. No. 5,283,173; Fields and Song, *Nature*, 340:245, 35 1989; Le Douarin et al., *Nucleic Acids Research*, 23:876,

- 37 -

1995; Vidal et al., *Proc. Natl. Acad. Sci. USA*, 93:10315-10320, 1996; and White, *Proc. Natl. Acad. Sci. USA*, 93:10001-10003, 1996). Generally, the two-hybrid methods involve *in vivo* reconstitution of two separable domains of a transcription factor. One fusion protein contains a *ynes* polypeptide fused to either a transactivator domain or DNA binding domain of a transcription factor (e.g., of Gal4). The other fusion protein contains a test polypeptide fused to either the DNA binding domain or a transactivator domain of a transcription factor. Once brought together in a single cell (e.g., a yeast cell or mammalian cell), one of the fusion proteins contains the transactivator domain and the other fusion protein contains the DNA binding domain. Therefore, binding of the *ynes* polypeptide to the test polypeptide (i.e., candidate antibacterial agent) reconstitutes the transcription factor. Reconstitution of the transcription factor can be detected by detecting expression of a gene (i.e., a reporter gene) that is operably linked to a DNA sequence that is bound by the DNA binding domain of the transcription factor. Kits for practicing various two-hybrid methods are commercially available (e.g., from Clontech; Palo Alto, CA).

The methods described above can be used for high throughput screening of numerous test compounds to identify candidate antibacterial (or anti-bacterial) agents. Having identified a test compound as a candidate antibacterial agent, the candidate antibacterial agent can be further tested for inhibition of bacterial growth *in vitro* or *in vivo* (e.g., using an animal, e.g., rodent, model system) if desired. Using other, art-known variations of such methods, one can test the ability of a nucleic acid (e.g., DNA or RNA) used as the test compound to bind *ynes*.

- 38 -

In vitro, further testing can be accomplished by means known to those in the art such as an enzyme inhibition assay or a whole-cell bacterial growth inhibition assay. For example, an agar dilution assay
5 identifies a substance that inhibits bacterial growth. Microtiter plates are prepared with serial dilutions of the test compound, adding to the preparation a given amount of growth substrate, and providing a preparation of bacteria. Inhibition of bacterial growth is
10 determined, for example, by observing changes in optical densities of the bacterial cultures.

Inhibition of bacterial growth is demonstrated, for example, by comparing (in the presence and absence of a test compound) the rate of growth or the absolute
15 growth of bacterial cells. Inhibition includes a reduction of one of the above measurements by at least 20%. Particularly potent test compounds may further reduce the growth rate (e.g., by at least 25%, 30%, 40%, 50%, 75%, 80%, or 90%).

20 Rodent (e.g., murine) and rabbit animal models of bacterial infections are known to those of skill in the art, and such animal model systems are accepted for screening antibacterial agents as an indication of their therapeutic efficacy in human patients. In a typical in
25 vivo assay, an animal is infected with a pathogenic strain of bacteria, e.g., by inhalation of bacteria such as *Streptococcus pneumoniae*, and conventional methods and criteria are used to diagnose the mammal as being afflicted with a bacterial infection. The candidate
30 antibacterial agent then is administered to the mammal at a dosage of 1-100 mg/kg of body weight, and the mammal is monitored for signs of amelioration of disease. Alternatively, the test compound can be administered to the mammal prior to infecting the mammal with the
35 bacteria, and the ability of the treated mammal to resist

- 39 -

infection is measured. Of course, the results obtained in the presence of the test compound should be compared with results in control animals, which are not treated with the test compound. Administration of candidate
5 antibacterial agents to the mammal can be carried out as described below, for example.

Pharmaceutical Formulations

Treatment includes administering a pharmaceutically effective amount of a composition
10 containing an antibacterial agent to a subject in need of such treatment, thereby inhibiting bacterial growth in the subject. Such a composition typically contains from about 0.1 to 90% by weight (such as 1 to 20% or 1 to 10%) of an antibacterial agent of the invention in a
15 pharmaceutically acceptable carrier.

Solid formulations of the compositions for oral administration may contain suitable carriers or excipients, such as corn starch, gelatin, lactose, acacia, sucrose, microcrystalline cellulose, kaolin,
20 mannitol, dicalcium phosphate, calcium carbonate, sodium chloride, or alginic acid. Disintegrators that can be used include, without limitation, micro-crystalline cellulose, corn starch, sodium starch glycolate and alginic acid. Tablet binders that may be used include
25 acacia, methylcellulose, sodium carboxymethylcellulose, polyvinylpyrrolidone (Povidone), hydroxypropyl methylcellulose, sucrose, starch, and ethylcellulose. Lubricants that may be used include magnesium stearates, stearic acid, silicone fluid, talc, waxes, oils, and
30 colloidal silica.

Liquid formulations of the compositions for oral administration prepared in water or other aqueous vehicles may contain various suspending agents such as methylcellulose, alginates, tragacanth, pectin, kelgin,

- 40 -

carrageenan, acacia, polyvinylpyrrolidone, and polyvinyl alcohol. The liquid formulations may also include solutions, emulsions, syrups and elixirs containing, together with the active compound(s), wetting agents, 5 sweeteners, and coloring and flavoring agents. Various liquid and powder formulations can be prepared by conventional methods for inhalation into the lungs of the mammal to be treated.

Injectable formulations of the compositions may 10 contain various carriers such as vegetable oils, dimethylacetamide, dimethylformamide, ethyl lactate, ethyl carbonate, isopropyl myristate, ethanol, polyols (glycerol, propylene glycol, liquid polyethylene glycol, and the like). For intravenous injections, water soluble 15 versions of the compounds may be administered by the drip method, whereby a pharmaceutical formulation containing the antibacterial agent and a physiologically acceptable excipient is infused. Physiologically acceptable excipients may include, for example, 5% dextrose, 0.9% 20 saline, Ringer's solution or other suitable excipients. Intramuscular preparations, a sterile formulation of a suitable soluble salt form of the compounds can be dissolved and administered in a pharmaceutical excipient such as Water-for-Injection, 0.9% saline, or 5% glucose 25 solution. A suitable insoluble form of the compound may be prepared and administered as a suspension in an aqueous base or a pharmaceutically acceptable oil base, such as an ester of a long chain fatty acid, (e.g., ethyl oleate).

30 A topical semi-solid ointment formulation typically contains a concentration of the active ingredient from about 1 to 20%, e.g., 5 to 10% in a carrier such as a pharmaceutical cream base. Various formulations for topical use include drops, tinctures,

- 41 -

lotions, creams, solutions, and ointments containing the active ingredient and various supports and vehicles.

The optimal percentage of the antibacterial agent in each pharmaceutical formulation varies according to the formulation itself and the therapeutic effect desired in the specific pathologies and correlated therapeutic regimens. Appropriate dosages of the antibacterial agents can be readily determined by those of ordinary skill in the art of medicine by monitoring the mammal for signs of disease amelioration or inhibition, and increasing or decreasing the dosage and/or frequency of treatment as desired. The optimal amount of the antibacterial compound used for treatment of conditions caused by or contributed to by bacterial infection may depend upon the manner of administration, the age and the body weight of the subject, and the condition of the subject to be treated. Generally, the antibacterial compound is administered at a dosage of 1 to 100 mg/kg of body weight, and typically at a dosage of 1 to 10 mg/kg of body weight.

Antibodies

The ynes polypeptide (or antigenic fragments or analogs thereof) can be used to raise antibodies useful in the invention, and such polypeptides can be produced by recombinant or peptide synthetic techniques (see, e.g., *Solid Phase Peptide Synthesis*, supra; Ausubel et al., supra). In general, the polypeptides can be coupled to a carrier protein, such as keyhole limpet hemocyanin (KLH), as described in Ausubel et al., supra, mixed with an adjuvant, and injected into a host mammal. A "carrier" is a substance that confers stability on, and/or aids or enhances the transport or immunogenicity of, an associated molecule. Antibodies can be purified,

- 42 -

for example, by affinity chromatography methods in which the polypeptide antigen is immobilized on a resin.

In particular, various host animals can be immunized by injection of a polypeptide of interest.

5 Examples of suitable host animals include rabbits, mice, guinea pigs, and rats. Various adjuvants can be used to increase the immunological response, depending on the host species, including but not limited to Freund's (complete and incomplete adjuvant), adjuvant mineral gels
10 such as aluminum hydroxide, surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, dinitrophenol, BCG (bacille Calmette-Guerin) and *Corynebacterium parvum*. Polyclonal antibodies are heterogeneous populations of antibody
15 molecules derived from the sera of the immunized animals.

Antibodies useful in the invention include monoclonal antibodies, polyclonal antibodies, humanized or chimeric antibodies, single chain antibodies, Fab fragments, F(ab')₂ fragments, and molecules produced using
20 a Fab expression library.

Monoclonal antibodies (mAbs), which are homogeneous populations of antibodies to a particular antigen, can be prepared using a yneS polypeptide (or immunogenic fragment thereof) and standard hybridoma
25 technology (see, e.g., Kohler et al., *Nature*, 256:495, 1975; Kohler et al., *Eur. J. Immunol.*, 6:511, 1976; Kohler et al., *Eur. J. Immunol.*, 6:292, 1976; Hammerling et al., In Monoclonal Antibodies and T Cell Hybridomas, Elsevier, NY, 1981; Ausubel et al., supra).

30 In particular, monoclonal antibodies can be obtained by any technique that provides for the production of antibody molecules by continuous cell lines in culture, such as those described in Kohler et al., *Nature*, 256:495, 1975, and U.S. Patent No. 4,376,110; the
35 human B-cell hybridoma technique (Kosbor et al.,

- 43 -

Immunology Today, 4:72, 1983; Cole et al., *Proc. Natl. Acad. Sci. USA*, 80:2026, 1983); and the EBV-hybridoma technique (Cole et al., Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, Inc., pp. 77-96, 1983). Such
5 antibodies can be of any immunoglobulin class including IgG, IgM, IgE, IgA, IgD, and any subclass thereof. The hybridomas producing the mAbs specific for yneS can be cultivated in vitro or in vivo.

Once produced, polyclonal or monoclonal antibodies
10 are tested for specific recognition of a yneS polypeptide in an immunoassay, such as a Western blot or immunoprecipitation analysis using standard techniques, e.g., as described in Ausubel et al., supra. Antibodies that specifically bind to the yneS polypeptide, or
15 conservative variants thereof, are useful in the invention.

Preferably, antibodies are produced using fragments of the yneS polypeptides that appear likely to be antigenic, by criteria such as high frequency of
20 charged residues. In one specific example, such fragments are generated by standard techniques of PCR, and are then cloned into the pGEX expression vector (Ausubel et al., supra). Fusion proteins are expressed in *E. coli* and purified using a glutathione agarose
25 affinity matrix as described in Ausubel, et al., supra.

If desired, several (e.g., two or three) fusions can be generated for each protein, and each fusion can be injected into at least two rabbits. Antisera can be raised by injections in a series, typically including at
30 least three booster injections. Typically, the antisera is checked for its ability to immunoprecipitate a recombinant yneS polypeptide, or unrelated control proteins, such as glucocorticoid receptor, chloramphenicol acetyltransferase, or luciferase.

- 44 -

Techniques developed for the production of "chimeric antibodies" (Morrison et al., *Proc. Natl. Acad. Sci.*, 81:6851, 1984; Neuberger et al., *Nature*, 312:604, 1984; Takeda et al., *Nature*, 314:452, 1984) can be used
5 to splice the genes from a mouse antibody molecule of appropriate antigen specificity together with genes from a human antibody molecule of appropriate biological activity. A chimeric antibody is a molecule in which different portions are derived from different animal
10 species, such as those having a variable region derived from a murine mAb and a human immunoglobulin constant region.

Alternatively, techniques described for the production of single chain antibodies (U.S. Patents
15 4,946,778; 4,946,778; and 4,704,692) can be adapted to produce single chain antibodies against a yneS polypeptide. Single chain antibodies are formed by linking the heavy and light chain fragments of the Fv region via an amino acid bridge, resulting in a single
20 chain polypeptide.

Antibody fragments that recognize and bind to specific epitopes can be generated by known techniques. For example, such fragments can include but are not limited to F(ab'), fragments, which can be produced by
25 pepsin digestion of the antibody molecule, and Fab fragments, which can be generated by reducing the disulfide bridges of F(ab'), fragments. Alternatively, Fab expression libraries can be constructed (Huse et al., *Science*, 246:1275, 1989) to allow rapid and easy
30 identification of monoclonal Fab fragments with the desired specificity.

Polyclonal and monoclonal antibodies that specifically bind to a yneS polypeptide can be used, for example, to detect expression of a yneS gene in other
35 bacteria. For example, a yneS polypeptide can be readily

- 45 -

detected in conventional immunoassays of bacteria cells or extracts. Examples of suitable assays include, without limitation, Western blotting, ELISAs, radioimmune assays, and the like.

5 Other Embodiments

 The methods of the invention also can be practiced with various fragments, variants, analogs, and derivatives of the yneS polypeptide described above that retain one or more of the biological activities of the yneS polypeptide. Naturally-occurring and non-naturally-occurring variants are useful in the invention. Compared with the naturally-occurring yneS gene sequences depicted in Figs. 1 and 2, the nucleic acid sequences encoding variants may have a substitution, deletion, or addition of one or more nucleotides. The preferred variants retain a function of the yneS polypeptide, e.g., as determined in a complementation assay.

 It is to be understood that, while the invention has been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope of the following claims. For example, other art-known assays to detect interactions of test compounds with proteins, or to detect inhibition of bacterial growth also can be used with the yneS genes and gene products (including homologs and orthologs of the sequences disclosed herein).

- 46 -

What is claimed is:

1. A method for identifying an antibacterial agent, the method comprising:

(a) contacting an S-yneS polypeptide with a test compound; and

(b) detecting an interaction of the test compound with the S-yneS polypeptide, wherein an interaction indicates that the test compound is an antibacterial agent.

2. A method of claim 1, further comprising:

(c) determining whether the test compound inhibits growth of bacteria, relative to growth of bacteria cultured in the absence of a test compound that interacts with the polypeptide, wherein inhibition of growth indicates that the test compound is an antibacterial agent.

3. A method of claim 1, wherein:

the S-yneS polypeptide is provided as a fusion protein comprising the S-yneS polypeptide fused to (i) a transcription activation domain of a transcription factor or (ii) a DNA-binding domain of a transcription factor;

the test compound is a fusion protein comprising the polypeptide fused to (i) a transcription activation domain of a transcription factor or (ii) a DNA-binding domain of a transcription factor, to interact with the fusion protein; and

interaction of the test compound with the polypeptide is detected as reconstitution of a transcription factor.

- 47 -

4. A method for identifying an antibacterial agent, the method comprising:

- (a) contacting an S-ynes polypeptide with a test compound;
- (b) detecting a decrease in function of the polypeptide contacted with the test compound; and
- (c) determining whether the test compound inhibits growth of bacteria, relative to growth of bacteria cultured in the absence of the test compound, wherein inhibition of growth indicates that the test compound is an antibacterial agent.

5. A method for identifying an antibacterial agent, the method comprising:

- (a) contacting a nucleic acid encoding S-ynes with a test compound; and
- (b) detecting an interaction of the test compound with the nucleic acid, wherein an interaction indicates that the test compound is an antibacterial agent.

6. A method of claim 5, further comprising:

- (c) determining whether a test compound inhibits growth of bacteria, relative to growth of bacteria cultured in the absence of the test compound that interacts the nucleic acid, wherein inhibition of growth indicates that the test compound is an antibacterial agent.

7. A method for identifying an antibacterial agent, the method comprising:

- (a) contacting an ortholog of an S-ynes polypeptide with a test compound; and
- (b) detecting an interaction of the test compound with the ortholog, wherein an interaction indicates that the test compound is an antibacterial agent.

- 48 -

8. A method of claim 7, further comprising:

(c) determining whether a test compound that interacts with the ortholog inhibits growth of bacteria, relative to growth of bacteria cultured in the absence of the test compound, wherein inhibition of growth indicates that the test compound is an antibacterial agent.

9. A method of claim 7, wherein:

the ortholog is provided as a fusion protein comprising the ortholog fused to (i) a transcription activation domain of a transcription factor or (ii) a DNA-binding domain of a transcription factor;

the test compound is a fusion protein comprising a polypeptide fused to (i) a transcription activation domain of a transcription factor or (ii) a DNA-binding domain of a transcription factor, to interact with the ortholog; and

interaction of the test polypeptide of the test compound with the ortholog is detected as reconstitution of a transcription factor.

10. A method for identifying an antibacterial agent, the method comprising:

(a) contacting an ortholog of an S-ynes polypeptide with a test compound;

(b) detecting a decrease in function of the ortholog contacted by the test compound; and

(c) determining whether the test compound inhibits growth of bacteria, relative to growth of bacteria cultured in the absence of the test compound, wherein inhibition of growth indicates that the test compound is an antibacterial agent.

11. A method for identifying an antibacterial agent, the method comprising:

- 49 -

(a) contacting a nucleic acid encoding an ortholog of S-yneS; and

(b) detecting interaction of the test compound with the nucleic acid, wherein interaction indicates that the test compound is an antibacterial agent.

12. The method of claim 11, further comprising:

(c) determining whether the test compound inhibits growth of bacteria, relative to growth of bacteria cultured in the absence of the test compound, wherein inhibition of growth indicates that the test compound is an antibacterial agent.

13. The method of any one of claims 1 to 12, wherein the test compound is selected from the group consisting of polypeptides, ribonucleic acids, small molecules, and deoxyribonucleic acids.

14. A method of any one of claims 7 to 12, wherein the ortholog is B-yneS.

15. A method of any one of claims 1 to 12, wherein the polypeptide is derived from a non-pathogenic *Streptococcus* strain.

16. A method of any one of claims 1 to 12, wherein the polypeptide is derived from a pathogenic *Streptococcus* strain.

17. A method of any one of claims 7 to 12, wherein the ortholog is derived from a gram positive bacterium.

- 50 -

18. A method of any one of claims 7 to 12, wherein the ortholog is derived from a pathogenic bacterium.

19. A method of claim 1 or 7, wherein the test compound is immobilized on a substrate, and interaction of the test compound with the ortholog is detected as immobilization of the ortholog on the immobilized test compound.

20. Use of an inhibitor of the function of an S-yneS polypeptide in the preparation of a medicament for treating a *Streptococcus pneumoniae* infection in a mammal.

21. A use of claim 21, wherein the inhibitor is an antibody that specifically binds the S-yneS polypeptide.

22. A pharmaceutical formulation comprising an antibacterial agent identified by the method of any one of claims 1 to 12, and a pharmaceutically acceptable excipient.

23. Use of an antibacterial agent identified by the method of any one of claims 1 to 12 in the preparation of a medicament for treating a bacterial infection in an organism.

24. A use of claim 24, wherein the antibacterial agent is an antibody that specifically binds S-yneS or an ortholog thereof.

25. The use claim 24, wherein the organism is a human or rodent.

Fig. 1

STRPW_MPI_yoeS

SEQ ID NO: 1 1 ATGATTACATAGTTTTTAAATCTAGCCTATCTGCTGGGTTGGATTCCATCTGCTCTGATTGGACAACTATCTTTGAAATCAATCTACGGGAGC 100
TACTAATUTTTATCAAAATAATTAGGATGGGATAGAGGACCGAAGCTAAGGTAGACCGAGAGCTTAACCTGTTTCATAAGAAAGTTAGTTAGATGGGCTCG

SEQ ID NO: 11

SEQ ID NO: 2 1 M I T I V L L I L A Y L L G S I P S O L W I G Q V F F O I N L R E H 34

101 ATGGTCTGTAACACTGGAAAGCAACACCTTCGGCATTAGGTAAGAAAGCTGGTATGGCAACCTTTGTGATTGACTTTTTCAAGGAACCTTAAC 200
TACCAAGACCAATTGTGACCTTGCTGCTGTGTGGAAGGCTTAAATCCAATCTTTGGACCATACCTTGGAAACACTAACTGAAAAAGTTTCTTGGGATCG

35 G S O N T G T T N T F R I L O K K A G M A T F V I D F F K G T L A 67

201 AAGGCTGCTTGGATTATTTTCATCTACAGGCGTTTCTCTCATCTTTGGACTTTTGGCTGTTATGGGCCATACCTTCCCTATCTTTCGAGGATTT 300
TTGGAGGAGGCTAATAAAAAGTAGATUTTCGGCAAGAGGAGAGTAGAAACCTGAAACCGACATAGCGGTATGGAGGGATAGAAACCTCTTAA

68 T L L P I I F H L Q Q V S P L I F G L L A V I G H T F P I F A G F 100

301 AAAGGTGTAAGGCTGTGCAACCACTGCTGGAGTGATTTTGGATTTCGGGCTATCTTCTGCTCTACCTTGGGATTATCTTCTTGGAGCTCTCTATC 400
TTTCCACCATTCGGACAGGCTTGCTCAGACCTCACTAAAGCCTTAAAGCGGTATGAAGACAGAGATGGAAAGCTTATAGAGAAAGCTCGAGAGATAG

101 K G G K A V A T S A G V I F G P A P I F C L Y L A I I F F G A L Y L 134

401 TTGGAGTATGATTTCACTGCTAGTGTCACAGCATGCTTGGGCTGTTATGGGCTTCTGCTCTTCCACTTTTGGTTTATCTGAGTAAGTATGA 500
AACGTCATATAAGTGACAGATCAGAGTGCTGATAGCTAAGCGGACATAGCCCGAGAGAGAGAGGTGAAAAACCAAAATAGGACTCATTTGACT

135 G S H I S L S S V T A S I A A V I G V L L P P L P G F I L S H Y D 167

501 CTCTCTTCACTGCTATCTTACGACTTCTAGTTTGTATTATCTTCTGTAAGGACAATATAGCTGATATCAAAATAAAACTGAAAAATTTGCTC 600
GAGAGAGAGTAGGATAATAGATCTGTGAGGATCAACTAATAGTAAGCAATATCTGTTATATGAGGATAGTTTTTATTTTACTTTTAAAGGAG

168 S L P I A I I L A L A S L I I I R H K D N I A R I K N K T E N L V 200

601 CCTTGGGATGAACTTAACCATCAAGATCTTAAAAATAA 642
GGAACCCCTAAGTTGGATTGGTATGTTCTAGGATTTTTTATT

201 P W G L N L T R Q D P K K • 213

2 / 4

Fig. 2

BACSU_yne2

SEQ ID NO: 3 1 ATGTTAATTGCTTTATTTGATTTATTTGGCTACTTGTATAGGAGCATTCATCTGGCTTAATTGTGGGCAAGCTTGGCAAGGAATTGATATTCCGGAGC 100
TACATTAAGCAATAAATACTAATAAAGCGGATGAACATATCGTCTGAAGGTAGACGGAAATTAACACCGCTTGGAAAGGTTTCTTTAACTATAAGCCCTCG
SEQ ID NO: 12
SEQ ID NO: 4 1 MLIALLIILAYLIQSIPSGLIIVGKLAKEIDIREN 34

101 ACGGAAGCGCAACTTAGCGCTACCATTCATTCOSTACATTTGGGTGTAAAGCTGTTGGGTCTCATAGCCGGAGATATTTTGAAAGGACACTGGC 200
TGGCTTGGCCGTTGATTCGGGATGGTTAGCTAAGGCATGTAACCCACATTTTGGACCAAGCCAGCATATCGGCTCTATAAACTTTCCCTGTGACCG
35 GSGH LGATNAFR T L G V K A G S V V I A G D I L K O T L A 67

201 AACTGCATTGCTTTTCTCATGCAATTTGATATTCAACCGCTTCTTGCAGGAGTCTTTGGGCTTTAGGCCAGGTGTTTCCCATCTTGGCAAAATTTAA 300
TTGACGTAAACGGAAGAGTACGTACACTATAAGTGGCGAAGAACGCTCTCGAAGCCGCAAAATCCGCTGCACAAAGGTTAGAAAGCGGTTTAAATTT
65 T A L P P L H S V D I E F L L A G V F A V L G E V F P I F A K F R 100

301 GCGGTAAAGCGGTGGGACATCAAGAGGCTTTTGTCTATTTTACCCACCCCTGTTATTTATCAGCAGGTTGGGATTTCTTCTATCTTTTATCTCA 400
CGGCCATTTGGCCACCGCTGTAGTCTCTCCGCAAAAGATAAATGCTGGGACAAATAAGTCTACCAAGCCCATAGAAGTAGAAAAATATGAAT
101 G G K A V A T S G G V L L P Y A P L L P I T M V A V F F I F L Y L T 134

401 CTAAATTTTCTCTCTCATGATTTTAAAGGATCTACTGTTTATATAAGTTTCTTGTCCATGATACGTTATTTGATTTGTTGTTACCTGCT 500
GATTTAAACAAAGAGAGTAGCTACAAATTTGCTAGATATGACATATATCAAGAAACAGGTACTATGCATAAATACTAACGCAATGGGACGA
135 K F V S L S S M L T Q I Y T V I Y S F F V H D T Y L L I V V T L L 167

501 CACTATTTTGTGATATACAGACCGGAGGCAATTAAGCAATTTATCAATAAAGCAAGCTTAAGTAAATGTTTATA 582
GTGATAAAGCACTATATGCTGTGGCTGCTGTAAATTTGCTTAATAGTATTTTGTCTGATTTTCATTTTACCAATAT
168 T I F V I Y R R R A N I R I I N K T E P K V K W L • 193

+ 4y. B

Strategy for the targeted deletions of genes in *S. pneumoniae*



product



1. PCR:

a) 1 + 2

Strap. DNA

b) 3 + 4

Strap. DNA

2. PCR:


1 + 4

product 1a), 1b), erm gene

Fig. 4.



Non-polar gene knockouts in *S. pneumoniae*



SEQUENCE LISTING

<110> Millennium Pharmaceuticals, Inc.

<120> USE OF YNES, ESSENTIAL BACTERIAL GENES AND
POLYPEPTIDES

<130> 07334/090001

<140> US 09/163,445

<141> 1998-09-30

<150> US 60/070,116

<151> 1997-12-31

<160> 12

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 213

<212> PRT

<213> Streptococcus pneumoniae

<400> 1

Met	Ile	Thr	Ile	Val	Leu	Leu	Ile	Leu	Ala	Tyr	Leu	Leu	Gly	Ser	Ile	1	5	10	15
Pro	Ser	Gly	Leu	Trp	Ile	Gly	Gln	Val	Phe	Phe	Gln	Ile	Asn	Leu	Arg	20	25	30	
Glu	His	Gly	Ser	Gly	Asn	Thr	Gly	Thr	Thr	Asn	Thr	Phe	Arg	Ile	Leu	35	40	45	
Gly	Lys	Lys	Ala	Gly	Met	Ala	Thr	Phe	Val	Ile	Asp	Phe	Phe	Lys	Gly	50	55	60	
Thr	Leu	Ala	Thr	Leu	Leu	Pro	Ile	Ile	Phe	His	Leu	Gln	Gly	Val	Ser	65	70	75	80
Pro	Leu	Ile	Phe	Gly	Leu	Leu	Ala	Val	Ile	Gly	His	Thr	Phe	Pro	Ile	85	90	95	
Phe	Ala	Gly	Phe	Lys	Gly	Gly	Lys	Ala	Val	Ala	Thr	Ser	Ala	Gly	Val	100	105	110	
Ile	Phe	Gly	Phe	Ala	Pro	Ile	Phe	Cys	Leu	Tyr	Leu	Ala	Ile	Ile	Phe	115	120	125	
Phe	Gly	Ala	Leu	Tyr	Leu	Gly	Ser	Met	Ile	Ser	Leu	Ser	Ser	Val	Thr	130	135	140	
Ala	Ser	Ile	Ala	Ala	Val	Ile	Gly	Val	Leu	Leu	Phe	Pro	Leu	Phe	Gly	145	150	155	160
Phe	Ile	Leu	Ser	Asn	Tyr	Asp	Ser	Leu	Phe	Ile	Ala	Ile	Ile	Leu	Ala	165	170	175	
Leu	Ala	Ser	Leu	Ile	Ile	Ile	Arg	His	Lys	Asp	Asn	Ile	Ala	Arg	Ile	180	185	190	
Lys	Asn	Lys	Thr	Glu	Asn	Leu	Val	Pro	Trp	Gly	Leu	Asn	Leu	Thr	His	195	200	205	
Gln	Asp	Pro	Lys	Lys												210			

<210> 2
 <211> 642
 <212> DNA
 <213> Streptococcus pneumoniae

<220>
 <221> CDS
 <222> (1)...(639)

<400> 2
 atg att aca ata gtt tta tta atc cta gcc tat ctg ctg ggt tcg att
 48
 Met Ile Thr Ile Val Leu Leu Ile Leu Ala Tyr Leu Leu Gly Ser Ile
 1 5 10 15
 cca tct ggt ctc tgg att gga caa gta ttc ttt caa atc aat cta cgc
 96
 Pro Ser Gly Leu Trp Ile Gly Gln Val Phe Phe Gln Ile Asn Leu Arg
 20 25 30
 gag cat ggt tct ggt aac act gga acg acc aac acc ttc cgc att tta
 144
 Glu His Gly Ser Gly Asn Thr Gly Thr Thr Asn Thr Phe Arg Ile Leu
 35 40 45
 ggt aag aaa gct ggt atg gca acc ttt gtg att gac ttt ttc aaa gga
 192
 Gly Lys Lys Ala Gly Met Ala Thr Phe Val Ile Asp Phe Phe Lys Gly
 50 55 60
 acc cta gca acg ctg ctt ccg att att ttt cat cta caa ggc gtt tct
 240
 Thr Leu Ala Thr Leu Leu Pro Ile Ile Phe His Leu Gln Gly Val Ser
 65 70 75 80
 cct ctc atc ttt gga ctt ttg gct gtt atc ggc cat acc ttc cct atc
 288
 Pro Leu Ile Phe Gly Leu Leu Ala Val Ile Gly His Thr Phe Pro Ile
 85 90 95
 ttt gca gga ttt aaa ggt ggt aag gct gtc gca acc agt gct gga gtg
 336
 Phe Ala Gly Phe Lys Gly Gly Lys Ala Val Ala Thr Ser Ala Gly Val
 100 105 110
 att ttc gga ttt gcg cct atc ttc tgt ctc tac ctt gcg att atc ttc
 384
 Ile Phe Gly Phe Ala Pro Ile Phe Cys Leu Tyr Leu Ala Ile Ile Phe
 115 120 125
 ttt gga gct ctc tat ctt ggc agt atg att tca ctg tct agt gtc aca
 432
 Phe Gly Ala Leu Tyr Leu Gly Ser Met Ile Ser Leu Ser Ser Val Thr
 130 135 140

gca tgc att gcg gct gtt atc ³ggg gtt ctg ctc ttt cca ctt ttt ggt
 480
 Ala Ser Ile Ala Ala Val Ile Gly Val Leu Leu Phe Pro Leu Phe Gly
 145 150 155 160
 ttt atc ctg agt aac tat gac tct ctc ttc atc gct att atc tta gca
 528
 Phe Ile Leu Ser Asn Tyr Asp Ser Leu Phe Ile Ala Ile Ile Leu Ala
 165 170 175
 ctt gct agt ttg att atc att cgt cat aag gac aat ata gct cgt atc
 576
 Leu Ala Ser Leu Ile Ile Ile Arg His Lys Asp Asn Ile Ala Arg Ile
 180 185 190
 aaa aat aaa act gaa aat ttg gtc cct tgg gga ttg aac cta acc cat
 624
 Lys Asn Lys Thr Glu Asn Leu Val Pro Trp Gly Leu Asn Leu Thr His
 195 200 205
 caa gat cct aaa aaa taa
 642
 Gln Asp Pro Lys Lys
 210

<210> 3
 <211> 193
 <212> PRT
 <213> Streptococcus pneumoniae

<400> 3
 Met Leu Ile Ala Leu Leu Ile Ile Leu Ala Tyr Leu Ile Gly Ser Ile
 1 5 10 15
 Pro Ser Gly Leu Ile Val Gly Lys Leu Ala Lys Gly Ile Asp Ile Arg
 20 25 30
 Glu His Gly Ser Gly Asn Leu Gly Ala Thr Asn Ala Phe Arg Thr Leu
 35 40 45
 Gly Val Lys Ala Gly Ser Val Val Ile Ala Gly Asp Ile Leu Lys Gly
 50 55 60
 Thr Leu Ala Thr Ala Leu Pro Phe Leu Met His Val Asp Ile His Pro
 65 70 75 80
 Leu Leu Ala Gly Val Phe Ala Val Leu Gly His Val Phe Pro Ile Phe
 85 90 95
 Ala Lys Phe Lys Gly Gly Lys Ala Val Ala Thr Ser Gly Gly Val Leu
 100 105 110
 Leu Phe Tyr Ala Pro Leu Leu Phe Ile Thr Met Val Ala Val Phe Phe
 115 120 125
 Ile Phe Leu Tyr Leu Thr Lys Phe Val Ser Leu Ser Ser Met Leu Thr
 130 135 140
 Gly Ile Tyr Thr Val Ile Tyr Ser Phe Phe Val His Asp Thr Tyr Leu
 145 150 155 160
 Leu Ile Val Val Thr Leu Leu Thr Ile Phe Val Ile Tyr Arg His Arg
 165 170 175

Ala Asn Ile Lys Arg Ile Ile Asn⁴ Lys Thr Glu Pro Lys Val Lys Trp
 180 185 190
 Leu

<210> 4
 <211> 582
 <212> DNA
 <213> Streptococcus pneumoniae

<220>
 <221> CDS
 <222> (1)...(579)

<400> 4
 atg tta att gct tta ttg att att ttg gcc tac ttg ata ggc agc att
 48
 Met Leu Ile Ala Leu Leu Ile Ile Leu Ala Tyr Leu Ile Gly Ser Ile
 1 5 10 15
 cca tct ggc tta att gtg ggc aag ctt gcc aaa gga att gat att cgg
 96
 Pro Ser Gly Leu Ile Val Gly Lys Leu Ala Lys Gly Ile Asp Ile Arg
 20 25 30
 gag cac gga agc ggc aac tta ggc gct acc aat gca ttc cgt aca ttg
 144
 Glu His Gly Ser Gly Asn Leu Gly Ala Thr Asn Ala Phe Arg Thr Leu
 35 40 45
 ggt gta aaa gct ggt tcg gtc gtc ata gcc gga gat att ttg aaa ggg
 192
 Gly Val Lys Ala Gly Ser Val Val Ile Ala Gly Asp Ile Leu Lys Gly
 50 55 60
 aca ctg gca act gca ttg cct ttt ctc atg cat gtt gat att cac ccg
 240
 Thr Leu Ala Thr Ala Leu Pro Phe Leu Met His Val Asp Ile His Pro
 65 70 75 80
 ctt ctt gca gga gtc ttt gcg gtt tta ggc cac gtg ttt ccc atc ttc
 288
 Leu Leu Ala Gly Val Phe Ala Val Leu Gly His Val Phe Pro Ile Phe
 85 90 95
 gcc aaa ttt aaa ggc ggt aaa gcc gtg gcg aca tca gga ggc gtt ttg
 336
 Ala Lys Phe Lys Gly Gly Lys Ala Val Ala Thr Ser Gly Gly Val Leu
 100 105 110
 cta ttt tac gca ccc ctg tta ttt atc acg atg gtt gcg gta ttc ttc
 384
 Leu Phe Tyr Ala Pro Leu Leu Phe Ile Thr Met Val Ala Val Phe Phe
 115 120 125

atc ttt tta tac ttg act aaa ttt⁵ gtt tct ctc tca tcg atg tta aca
432
Ile Phe Leu Tyr Leu Thr Lys Phe Val Ser Leu Ser Ser Met Leu Thr
130 135 140

ggg atc tat act gtt ata tat agt ttc ttt gtc cat gat acg tat tta
480
Gly Ile Tyr Thr Val Ile Tyr Ser Phe Phe Val His Asp Thr Tyr Leu
145 150 155 160

ttg att gtc gtt acc ctg ctc act att ttt gtg ata tac aga cac cga
528
Leu Ile Val Val Thr Leu Leu Thr Ile Phe Val Ile Tyr Arg His Arg
165 170 175

gcg aac att aaa cga att atc aat aaa aca gaa cct aaa gta aaa tgg
576
Ala Asn Ile Lys Arg Ile Ile Asn Lys Thr Glu Pro Lys Val Lys Trp
180 185 190

tta taa
582
Leu

<210> 5
<211> 21
<212> DNA
<213> B. subsilis

<400> 5
gtgttcgtgc tgacttgac c
21

<210> 6
<211> 21
<212> DNA
<213> B. subtilis

<400> 6
gaattatttc ctcccgttaa a
21

<210> 7
<211> 21
<212> DNA
<213> Steptococcus pneumoniae

<400> 7
accactccg tgaagtccac c
21

<210> 8

6

<211> 21
 <212> DNA
 <213> Streptococcus pneumoniae

<400> 8
 gaacccagca gataggctag g
 21

<210> 9
 <211> 21
 <212> DNA
 <213> Streptococcus pneumoniae

<400> 9
 gaacctaacc catcaagatc c
 21

<210> 10
 <211> 24
 <212> DNA
 <213> Streptococcus pneumoniae

<400> 10
 cagttttaga tggctttaac agcg
 24

<210> 11
 <211> 642
 <212> DNA
 <213> Streptococcus pneumoniae

<400> 11
 tactaatgtt atcaaaataa ttaggatcgg atagacgacc caagctaagg
 tagaccagag 60
 acctaacctg ttcataagaa agtttagtta gatgcgctcg taccaagacc
 attgtgacct 120
 tgctgggttg ggaaggcgta aaatccattc tttcgaccat accgttggaa
 acactaactg 180
 aaaaagtttc cttgggatcg ttgcgacgaa ggctaataaa aagtagatgt
 tccgcaaaga 240
 ggagagtaga aacctgaaaa ccgacaatag ccggtatgga agggatagaa
 acgtcctaaa 300
 tttccaccat tccgacagcg ttggtcacga cctcactaaa agcctaaacg
 cggatagaag 360
 acagagatgg aacgctaata gaagaaacct cgagagatag aaccgtcata
 ctaaagtgc 420
 agatcacagt gtcgtagcta acgccgacaa tagccccaag acgagaaagg
 tgaaaaacca 480
 aataggact cattgatact gagagagaag tagcgataat agaatcgtga
 acgatcaaac 540
 taatagtaag cagtattcct gttatatcga gcatagtttt tattttgact
 tttaaaccag 600
 ggaacccta acttggattg ggtagttcta ggatttttta tt
 642

<210> 12
<211> 582
<212> DNA
<213> *Streptococcus pneumoniae*

<400> 12
tacaattaac gaaataacta ataaaaccgg atgaactatc cgtcgtaagg
tagaccgaat 60
taacacccgt tcgaacgggt tccttaacta taagccctcg tgccttcgcc
gttgaatccg 120
cgatgggttac gtaaggcatg taaccacacat tttcgaccaa gccagcagta
tcggcctcta 180
taaaactttc cctgtgaccg ttgacgtaac ggaaaagagt acgtacaact
ataagtgggc 240
gaagaacgtc ctcagaaacg ccaaaatccg gtgcacaaag ggtagaagcg
gtttaaattt 300
ccgccatttc ggcaccgctg tagtcctccg caaaacgata aaatgcgtgg
ggacaataaa 360
tagtgctacc aacgccataa gaagtagaaa aatatgaact gatttaaaca
aagagagagt 420
agctacaatt gtccttagat atgacaatat atatcaaaga aacaggtact
atgcataaat 480
aactaacagc aatgggacga gtgataaaaa cactatatgt ctgtggctcg
cttgtaattt 540
gcttaatagt tattttgtct tggatttcac tttaccaata tt
582

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/22665

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C1201/18 G01N33/94 C07K14/315 C07K14/32

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12Q G01N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A,P	WO 99 33871 A (MILLENNIUM PHARMACEUTICALS INCORPORATED) 8 July 1999 (1999-07-08) claim 22; figures 1-20	1-26
A	CHEMICAL ABSTRACTS, vol. 128, no. 13, 30 March 1998 (1998-03-30) Columbus, Ohio, US; abstract no. 150233, XP002128453 abstract & F. KUNST ET AL.: "The complete genome sequence of the gram-positive bacterium Bacillus subtilis" NATURE (LONDON), vol. 390, no. 6657, 1997, pages 249-256, London UK	1-26

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

21 January 2000

Date of mailing of the international search report

03/02/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 851 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bohemen, C

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/22665

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9933871 A	08-07-1999	AU 2024399 A	19-07-1999